

THE UNITED STATES OF AMERICA

To am to vnom these presents sham come: Irito-Unp Aorth America, Inc.

MOCCUS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE DIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT ED BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

POTATO

'FL 1900'

In Jestimone Mercent, I have hereunto set my hand and caused the seal of the Hunt Harrety Frotection Office to be affixed at the City of Washington, D.C. this sixth day of February, in the year two thousand and seven.

Atlast:

Olmbe

Commissioner Plant Variety Protection Office Agricultural Marketing Service a government of

Agriculture

Secretary, Recot, Inc.

SET-470 (2-98) designed by the Plant Variety Protection Office with WordPerfect 6.0a, Replaces STD-470 (6-98) which is obsolete.

(See reverse for instructions and information collection builden statement)

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvpindex.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

ITEM

- 19a. Give:
- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97,103).
- See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

NIA

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filling a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

EXHIBIT A. Origin and Breeding History of the Variety


The variety FL 1900 originated in the Frito-Lay, Inc., private potato breeding program. In 1990, a cross was made by Robert W. Hoopes at the Frito-Lay Research facility near Rhinelander, Wisconsin, between the varieties FL 1700 and S440. The variety FL 1700 was chosen as a breeding parent because of its excellent potato chip color, high solids, regular round tubers of the right size for chipping, resistance to common scab, and resistance to the "golden" cyst nematode, *Globodera rostochiensis* race RO1. S440, a University of Wisconsin breeding line, was chosen as a parent because of its desirable chip color, high solids content, and good tuber shape and size.

Seeds from the cross FL 1700 x S440 were sown in the greenhouse near Rhinelander in the summer of 1990 and the resulting tubers harvested in late fall of that year. Seedling tubers were planted in the field in the spring of 1991. One of the selections from this progeny was given the designation "RD 78-91-5." This selection was tested for four years in the fields in Rhinelander, with solids measurements taken and potato chip samples made after each harvest. RD 78-91-5 was found to have consistently good tuber type for shipping, high solids (usually higher than the variety 'Atlantic'), and good chip quality.

In 1995, RD 78-91-5 was given the number "FL 1900" and underwent experimental testing in the national variety trials conducted by Frito-Lay in Florida, Texas, California, New Mexico, North Dakota, Wisconsin, Michigan, and Maine. It has since been tested in the Advanced Clonal Evaluation program and found to be well-adapted to growing conditions for the spring crop in the Southeast, producing good yields, solids higher than those of Atlantic, lower internal defects than Atlantic, and good chip quality. FL 1900 has been tested by Dr. Bill Brodie of Cornell University in Ithaca, New York, and found to be resistant to the "golden" potato cyst nematode *Globodera rostochiensis* race RO1.

The variety FL 1900 has been uniform and stable since its origin as a single plant in 1991. No variants of FL 1900 have been observed.

Tissue culture plantlets of FL 1900 were established beginning in 1994 and are maintained at the Frito-Lay Research facility near Rhinelander, Wisconsin, with a duplicate maintained by the University of Wisconsin Biotron facility in Madison.



Trial Data for 'FL 1900'

Location	Yield	Solids
Davis, California	458	20.2
Holyoke, Colorado	278	17.1
Rochester, Indiana	205	18.2
Garden City, Kansas	356	18.9
Rhodesdale, Maryland	529	18.0
Cribou, Maine	264	20.1
Three Rivers, Michigan	414	18.5
Charleston, Missouri	200	16.3
Plymouth, North Carolina	252	15.9
Clovis, New Mexico	243	16.8
Clovis, New Mexico	264	16.3
Clovis, New Mexico	328	17.4
Hermiston, Oregon	438	18.5
Antigo, Wisconsin	443	19.3
Hancock, Wisconsin	414	18.9

The data indicated above is from observation plots conducted by Frito-Lay in 1996. Frito-Lay North America, Inc. does not participate in the National Trial Test.

EXHIBIT B. Statement of Distinctness

As a chipping variety, FL 1900 is most similar to Atlantic. FL 1900 differs from Atlantic in that the corolla color of FL 1900 is a dark violet, RHS N88B, while the flower color of Atlantic is a light violet, RHS N82D. FL 1900 has more prominent white tips, whiter than RHS 155D, on the corolla and more prominent white stripes, whiter than RHS 155D, on the outer surface of the corolla extending from the tips towards the center, while Atlantic has RHS 155B white tips on the and RHS 155B white stripes on the corolla. Additionally, the stem anthocyanin of FL 1900 is a dark violet, RHS 86A to RHS 86B, than the stem anthocyanin of Atlantic, a faint violet of RHS N78B to RHS N78C.

The isozyme pattern of FL 1900, as established by Dr. David Douches of Michigan State University, is unique among known North American varieties. This is detailed in Exhibit D, Additional Description.

Comparison of Inner Surface of Corolla of 'FL 1900' with 'Atlantic'



'FL 1900'



'Atlantic'

NAME OF APPLICANT(S)			FOR OFFICIAL USE OF	VLY
Recot, Inc.			PVPO NUMBER 20000026	
ADDRESS (Street and No. or R.	.F.D. No., City, State, and Zip C	Code)	VARIETY (V) NAME	
5000 Hopyard Drive			FL1900	•
Suite 460 Pleasanton, CA 9458	8		TEMPORARY OR EXPERIM	ENTA
ricasancon, on 5450			DESIGNATION	
			RD 78-91-5	
		·		
REFERENCE VAR	IETIES: Enter the ref	erence variety na	me in the appropriate	box
Reference Variety 1 (R1)	Reference Variety 2 (R2)	Reference Variety 3	(R3) Reference Variety	(R4)
Atlantic		·		•
. MARKET CHARACTERIST	TCS:	<u> </u>		· · ·
	. ACO.			
MARKET CLASS:		e, e		
1 = Yellow-flesh tablest5 = Russet tablestock;	ock; 2 = Round-white tablesto 6 = Other	ck; 3 = Chip-processing	g; 4 = Frozen-processing;	
T AND THE MORE THAN				
V 3	R1 3 R2	R3	R4	
. PLANT CHARACTERISTIC	CS:			
GROWTH HABIT: (See) $3 = \text{Erect} (>45^{\circ} \text{ with g})$	figure 1) ground); 5 = Semi-erect (30-45	o with ground); 7 = Spr	reading.	
V 5	R1 5 R2	R3	R4	
				Y
TYPE: 1 = Stem (foliage open,	, stems clearly visible); $2 = Int$	ermediate; 3 = Leaf (F	oliage closed, stems hardly visib	le)
				\neg
V 2	R1 R2	R3	R4	
	<u> </u>			
MATURITY: Days after I	planting (DAP) at vine senescen	ce		
V 105 R	1 100 R2	R3	R4	
PLANTING DATE:			D4	
Mid-April 1997 R1 A	pril 1997 R2	R3	R4	
REGION/AREA:	ma beatenes of color			
		11 1	الحا	4
Maryland• R1	R2	R3	R4	
Maryland• R1	orth East R2 C per letter 8/15/20	R3	R4	

MAT	URITY	CL	LSS:
_		_	

1 = Very Early (<100 DAP); 2 = Early (100-110 DAP); 3 = Mid-season (111-120 DAP); 4 = Late (121-130 DAP); 5 = Very Late (>130 DAP).

v 2-3

R1 2

R2

R3

R4

3. STEM CHARACTERISTICS: Measure at early first bloom

*

STEM ANTHOCYANIN COLORATION:

1 = Absent; 3 = Weak; 5 = Medium; 7 = Strong; 9 = Very Strong

V 7

R1 5

R2

R3

R4

STEM WINGS: (See figure 12)

1 = Absent; 3 = Weak; 5 = Medium; 7 = Strong; 9 = Very Strong

v 5

R1 5

R2

R3

R4

4. LEAF CHARACTERISTICS:

LEAF COLOR: (Observe fully developed leaves located on middle 1/3 of plant)

1 = Yellowish-green; 2 = Olive-green; 3 = Medium green; 4 = Dark green; 5 = Grey-green; 6 = Other_

V 2

R1 2

R2

R3

R4

LEAF COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Chart

(Observe fully developed leaves located on middle 1/3 of plant & circle the appropriate color chart)

V 137A

R1 137A

R2

R3

R4

LEAF PUBESCENCE DENSITY:

1 = Absent; 2 = Sparse; 3 = Medium; 4 = Thick; 5 = Heavy

V 5

R1 7

R2

R3

R4

LEAF PUBESCENCE LENGTH:

1 = None; 2 = Short; 3 = Medium; 4 = Long; 5 = Very long

V

R1

R2

R3

R4

(Note: Descriptor #19 can be used to describe the type and length of the glandular trichomes observed.)

*

LEAF SILHOUETTE: (See figure 2)

1 = Closed; 3 = Medium; 5 = Open

V 5

R1 5

R2

R3

R4

· .·		OLES AN = Absent;					Strong;	9 = Ve	ry Strong	3					
	v	3		R1	1		R2			R3			R4		
	1 =	STIPULI = Absent;		nall; 5 =		m; 7 =			· .						
	V			R1			R2		<u> </u>	R3			R4		
	1 =	MNAL LE Narrowl Obovate;	y ovate;	2 = M	edium ov	ate; 3 =		ly ovate;	4 = La	nceolate;	5 = Elli	ptical;			
	V	2		R1			R2			R3			R4		
		INAL LE Acute; 2							= Other_						
٠.	V	3		R1			R2]	R3			R4		
*	TERM	INAL LE						•	uncate;	6 = Lob	ed; 7 =	Other			
	V	4		R1	4		R2			R3		·	R4		
*		NAL LEA					dium; 5	= Stron	g						
	V	4		R1	3		R2			R3			R4		
		ER OF PR RAGE:	RIMARY	/ LEAFI	LET PAI	RS: (Se	e figure	11)							
	V	3		R1			R2			R3			R4		
RAN	VGE:											·			·
V	t	0	R1	t	0	R2		to	R	3	to	R	1	to	
		RY LEAF Acute; 2						use; 5 =	Other_			<u> </u>			
	V	3	•	R1		•	R2			R3			R4		

PRIMARY LEAFLET SIZE: 1 = Very Small; 2 = Small; 3 = Medium; 4 = Large; 5 = Very Large	
V 3 R1 3 R2 R3 R4	
PRIMARY LEAFLET SHAPE: (See figure 3 & 11) 1 = Narrowly ovate; 2 = Medium ovate; 3 = Broadly ovate; 4 = Lanceolate; 5 = Elliptical; 6 = Obovate; 7 = Oblong; 8 = Other	
V 2 R1 R2 R3 R4	
PRIMARY LEAFLET BASE SHAPE: (See figure 5 & 11) 1 = Cuneate; 2 = Acute; 3 = Obtuse; 4 = Cordate; 5 = Truncate; 6 = Lobed; 7 = Other	
V 4 R1 R2 R3 R4	
NUMBER OF SECONDARY AND TERTIARY LEAFLET PAIRS: (See figure 11) AVERAGE:	
V 10 R1 R2 R3 R4	
RANGE: V 8 to 10 R1 to R2 to R3 to R4 to	-
5. INFLORESCENCE CHARACTERISTICS:	
NUMBER OF INFLORESCENCE / PLANT: AVERAGE:	
V R1 R2 R3 R4	
RANGE: V to R1 to R2 to R3 to R4 to	-
NUMBER OF FLORETS / INFLORESCENCE:	
AVERAGE: V R1 R2 R3 R4	
RANGE:	.
V to R1 to R2 to R3 to R4 to	
*COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Cha (Measure predominant color of newly open flower & circle the appropriate color chart)	urt
V 88B R1 82D R2 R3 R4	

COROLLA INNER SURFACE COLOR: (Measure predomin 1 = White; 2 = Red-violet; 3 = Blue-violet; 4 = Other	
V 4 R1 4 R2	R3 R4
COROLLA SHAPE: (See figure 6) 1 = Very rotate; 2 = Rotate; 3 = Pentagonal; 4 = Sem	i-stellate; 5 = Stellate
V 3 R1 4 R2	R3 R4
CALYX ANTHOCYANIN COLORATION: 1 = Absent; 3 = Weak; 5 = Medium; 7 = Strong; 9 =	Very strong
V 5 R1 1 R2	R3 R4
ANTHER COLOR CHART VALUE: Royal Horticulture Soc (Measure when newly opened flower is fully expanded and circle in	ciety Color Chart or Munsell Color Chart the appropriate color chart)
V 14A R1 2A R2	R3 R4
ANTHER SHAPE: (See figure 7) 1 = Broad cone; 2 = Narrow cone; 3 = Pear shape cone	; 4 = Loose; 5 = Other
V 2 R1 3 R2	R3 R4
POLLEN PRODUCTION: 1 = None; 3 = Some; 5 = Abundant	
V 5 R1 3 R2	R3 R4
STIGMA SHAPE: (See figure 8) 1 = Capitate; 2 = Clavate; 3 = Bi-lobed	
V 1 R1 R2	R3 R4
STIGMA COLOR CHART VALUE: Royal Horticulture Socie (Circle the appropriate color chart)	ety Color Chart or Munsell Color Chart
137A R1 R2	R3 R4
BERRY PRODUCTION: (Under field conditions) 1 = None; 3 = Low; 5 = Moderate; 7 = Heavy; 9 = Ver	ry heavy
V 5 P1 5 P2	D2 D4

21D-4/6 (01-86)

Page 7 of 19

TUBER CHARACT	ERISTICS:			
* PREDOMINANT	SKIN COLOR:	· :		
			Tan; $6 = Brown$; $7 = 1$	Pink; 8 = Red;
9 = Purplish-r	ed; 10 = Purple; 11 =	Dark purple-black; 12	z = Other	
V 6	R1 6	R2	R3	R4
DREDOMINANT	SKIN COLOR CHART	WALLE David Hoti	aultura Sariety Calar Ch	ent on Managell Calas Ch. 4
(Circle the appropri		VALUE: Royal Horus	culture Society Color Ch	art or Munsell Color Chart
			The last	
V 199B	R1	R2	R3	R4
	****		·	
SECONDARY SKI	IN COLOR:			
1 = Absent; 2	= Present, please descri	be:		
		7 [
V	R1	R2	R3	R4
SECONDARY SKI	N COLOR CHART VA	LUE: Royal Horticult	ure Society Color Chart	or Munsell Color Chart
(Circle the approprie	·	• • • • • • • • • • • • • • • • • • •		
v	R1	R2	R3	R4
		J [
SECONDARY SKT	N COLOR DISTRIBUT	TON:		
			Spectacled; 6 = Stippl	ed; 7 = Other
v	R1	R2	R3	R4
<u> </u>	KI	. RZ	I NO	ICT .
SKIN TEXTURE:				
	= Rough (flaky); 3 = N	Vetted; 4 = Russetted;	5 = Heavily russetted;	6 = Other
***	Dr.	D2	D2	R4
V 3	R1	R2	R3	K4
TUBER SHAPE: (5	See figure 10); $3 = Ove$	$\mathbf{a} \cdot \mathbf{A} = \mathbf{Oblong} \cdot 5 = 1$	Long: 6 = Other	
T = Compresses	, 1 - 100110, 3 - 011	, - Oblong, 5	l	1
v 2-3	R1 3	R2	R3	R4
		e de la companya de La companya de la co		
	22			
TUBER THICKNES 1 = Round; 2 =	SS: : Medium thick; 3 = Sl	ightly flattened; 4 = 1	Flattened; 5 = Other_	
				1
1 37	701	l pa	D3	R4

			
TUBER LENGTH (mm):			
AVERAGE:			
v 77.5 R1 75	R2	R3	R4
RANGE:			
V 62 to 106 R1 51 to 115	R2 to	R3 to	R4 to
STANDARD DEVIATION:			
V 9.2 R1 14.8	R2	R3	R4
AVERAGE WEIGHT OF SAMPLE TAI	ŒN:		
V R1	R2	R3	R4
TUBER WIDTH (mm):			
AVERAGE:			
V 60.6 R1 64	R2	R3	R4
RANGE:			——————————————————————————————————————
V 44 to 75 R1 47 to 86	R2 to	R3 to	R4 to
STANDARD DEVIATION:			
V 5.2 R1 9.1	R2	R3	R4
AVERAGE WEIGHT OF SAMPLE TAKE V R1		Da	
V K I	[R2]	R3	R4
TUBER THICKNESS (mm):			
AVERAGE:	<u> </u>	· · · · · · · · · · · · · · · · · · ·	
V 45 R1 51	R2	R3	R4
RANGE:			
V 38 to 53 R1 35 to 66	R2 to	R3 to	R4 to
STANDARD DEVIATION:			
V 3.5 R1 6.7	R2	R3	R4
_AVERAGE WEIGHT OF SAMPLE TAKE	CN:		
V R1	R2	R3	R4
			<u> </u>
TUBER EYE DEPTH:			
1 = Protruding; 2 = Shallow; 3 = Intern	mediate; 4 = Deep; 5 =	Very deep	
V 2 R1 3	R2	R3	R4

Page 9 of 19

1 = Protruding	EYES; 2 = Shallow; 3 = Inter	mediate; 4 = Deep; 5 =	Very deep	
v	R1	R2	R3	R4
NUMBER EYE / T AVERAGE: V 5	TUBER:	R2	R3	R4
RANGE:				
V to	R1 to	R2 to	R3 to	R4 to
DISTRIBUTION Of 1 = Predominan	F TUBER EYES: tly apical; 2 = Evenly dis	tributed	R3	R4
	TUBER EYEBROWS: ent; 2 = Slight prominence R1 1	e; 3 = Medium promine	nce; 4 = Very prominent;	5 = Other
PRIMARY TUBE (Circle the appropriat) V 155A	the state of the s	RT VALUE: Royal Hort	iculture Society Color Char	t or Munsell Color Chart
SECONDARY TUBE 1 = Absent; 2 =	ER FLESH COLOR: Present, please describe:			
v	R1	R2	R3	R4
SECONDARY TUBE (Circle the appropriate		RT VALUE: Royal Hort	iculture Society Color Char	rt or Munsell Color Chart
NUMBER OF TUBE 1 = Low (<8); 2	RS / PLANT: = Medium (8 -15); 3 =	High (>15)		
V 2	R1 2 slightly lowers	R2	R3	R4

		•		RISTICS		**************************************	1 DEC	SCHOOL NEW		CORTO		a Horom		
	DISEA	SES KEA	CHON	0 = r $5 = N$	MODERA	ATELY	1 = RES SUSCEP	TIBLE;	7 = SUS	CEPTIB	LE; 9=I	RESISTA HGHLY	ANT; SUSCE	PTIBLI
	BACT	ERIAL I	ZING R	OT, FOI	LIAR RE	ACTIO	N:						ě	
	V	7		R1			R2.			R3			R4	
	BACT	ERIAL F	ING R	OT. TU	BER REA	- ACTIOI	V:		- 					
5 -	V	7		R1			R2			R3			R4	
	LATE	E BLIGH	T:						· .			- :		
	V	-9- 4		R1	7		R2			R3			R4	
	PLRV	4	ROLL):	· letter	8/15/2	2006			. ·			.	·	
	v	0		R1			R2			R3			R4	
	PVX:	L	nc ger	letter	8/15/	2006	• .		•			.		
	V			R1			R2		÷	R3			R4	
	PVY:		. '											
	v	9		R1		•	R2			R3	(4)		R4	
	ОТНЕ	R: Sof	t Ro	t			<u> </u>							
	v	6		R1	7-8	* .	R2			R3			R4	
•	ОТНЕ	R•	LMC	861	letter	8/15/	2006					•		
	v			R1			R2			R3			R4	
PE	STS CH	ARACTI	RISTI									· ·	· · · · · · · · · · · · · · · · · · ·	
			•		200000	- n	DOTOM 4 N	m a	MODE	D A FERRY N	z meoro	TD 4 N TOS.	· · · · · · · · · · · · · · · · · · ·	
,	resi ki	EACTIO					ESISTAN EPTIBLI						CEPTIBI	Æ
	GOLD	en nem	ATODE	C:										
	v	1		R1	:		R2	: :	•	R3			R4	
	OTHE	રઃ								,				
	V			R1		•	R2			R3			R4	
GE	NE TRA	ATS:	:					<u> </u>	· · : ·		•			· · ·
		ON OF	CENES	•			YES			П	NO	1.41		
					baad	- Office-		ion:		X			• .	٠.
— Li	LIES, C	escribe t	ne gene	(s) intro	nacea or	attaen	informat	7011:			. · · · · ·			<u>.</u>
		<u> </u>												
_					<u> </u>			·						

CTD_47K (01.04)

Page 11 of 19

8. DISEASES CHARACTERISTICS: (continued)	2200000268
POTATO VIRUS X (PVX)	
V O R1 R2 R3	R4
POTATO VIRUS Y (PVY)	
V C R1 R2 R3	R4
V O R1 R2 R3	R4
POTATO VIRUS A (PVA)	
V O R1 R2 R3	R4
GOLDEN NEMATODE (Globodera)	
V R1 R2 R3	R4
ROOT – KNOT NEMATODE (Meloidogyne)	
V O R1 R2 R3	R4
OTHER DISEASE	
V R1 R2 R3	R4
PHYSIOLOGICAL DISORDER 1 = Malformed shape	v heart 5 = Internal necrosis
V R1 R2 R3	R4
9. PESTS CHARACTERISTICS:	
PEST REACTION: 0 = Not Tested 1 = Highly Resistant 2 = Resistant Few Symptom 4 = Moderately Resistance 5 = Intermedia Susceptible 6 = Moder 7 = Susceptible 9 = Highly Susceptible	s 3 = Resistance Few Lessions in Number and Size rate Susceptible
COLORADO POTATO BEETLE (CPB) (Leptinotarsa)	
V R1 R2 R3	R4
GREEN PEACH APHID (Myzus)	
V R1 R2 R3	R4
	[]
V R1 R2 R3	R4
V R1 R2 R3	R4

	CHIEF MARKET			-			•					
	Processing to	nake po	tato chi	ps		· · · · · · · · · · · · · · · · · · ·						
	SPECIFIC GRAVI $1 < 1.060; 2 =$					1.080-1.089	; 5 > 1.	090				
			- i	7		·	· ·	 	· 		-	
	V 4	R	1].	R2			R3		R	4	
	TOTAL GLYCOAL	LKALOII	D CONTEN	T (mg.	/ 100 g.	fresh tuber)					
	V 8.2	R	1 9.2		R2			R3		R	4	
	corresponding proto	col.		· · · · · · · · · · · · · · · · · · ·	· .							
_									******			
									•			
	IEMICAL IDENTII Describe chemical tr	aits of the	candidate			n its identifi	cation (e	g. protei	n or DNA	electrop	ohoresis)	
		aits of the	candidate			n its identifi	cation (e	g. protei	n or DNA	electrop	phoresis)	•
	Describe chemical tr Please attach data ar	aits of the	candidate			n its identifi	cation (e	g. protei	n or DNA	electrop	phoresis)	
	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ol.	n its identifi	cation (e	g. protei	n or DNA	electrop	ohoresis)	
	Describe chemical tr Please attach data ar	aits of the	e candidate responding	protoco	ol.	n its identifi	cation (e	.g. protei	n or DNA	electrop	ohoresis)	
AI	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	ohoresis)	
1	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	ohoresis)	
AI.	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	ohoresis)	
\I	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
1	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
AI	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
AI	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
AI	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
AI	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
AL	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	phoresis)	
AL	Describe chemical tr Please attach data an See Exhibit D	aits of the	e candidate responding	protoco	ISTICS:					electrop	ohoresis)	


					Exhibit C (Pota
NAME OF APPLICANT (S)		TEMPORARY OR EXPERIMENTAL DESIGNA	TION	VARIETY NAMI	E
ADDRESS (Street and No. or RD No., Cit	ty, State, Zip Code, and Country)			FOR OFFICIAL	USEONLY
				PVPO NUMBER	3
		TOTAL POLICE AND			
REFERENCE VARIETIES: Ente	I		Defense Mediate	2 (D2)	Deference Verichy 4 (D4)
Application Variety (V)	Reference Variety 1 (R1)) Reference Variety 2 (R2)	Reference Variety	3 (K3)	Reference Variety 4 (R4)
					•
PLEASE READ ALL INSTRU	CTIONS CAREFULLY:				
1. MARKET CHARACTERISTIC			90	000	0 2 6 8
*MARKET CLASS:			5 (VUU	
	stock 2 = Round-white Ta 6 = Other	ablestock 3 = Chip-processing 4 =	Frozen-processing		
V	R1	R2 R3	R4		
		<u> </u>	i /\ i	ا م ما	
2. LIGHT SPROUT CHARACTE		Comparison Vo	riety= AT	ICCUTU	
*LIGHT SPROUT: GE 1 = Spherical 2 =		= Broad cylindrica 5 = Narrow o	ylindrical 6 = Other	Г	
X 2	D1 //	D2 D2			
V 3	R1 4	R2 R3	R4		
*LIGHT SPROUT BAS 1 = Absent 2 = We	E: PUBESCENCE OF TIP	= Strong 5 = Very Strong			
T - Absent 2 - VVC	eak 3 - Medium 4	- Strong 5 - Very Strong			
v 3	R1 3	R2 R3	R4		
#I ICUT EPPOUT PAG	E: ANTHOCYANIN COLO	DRATION			
1 = Green 2 = Red-		4 = Other(describe)			
V 3	R1 2	D2 D2	\Box		
V 5	$R1 \mathcal{L}$	R2 R3	R4		
*LIGHT SPROUT BAS 1 = Absent 2 = We		OCYANIN COLORATION (IF PRESE Strong 5 = Very Strong	NT)		
1 - Absent 2 - We	ak 5 – Iwedium 4 =	- Strong 5 = Very Strong			
v 5	R1 5	R2 R3	R4		
* 1 1017 2 2 2 2 2 2 2	LIADIT				
* LIGHT SPROUT TIP: 1 = Closed 2 = Int	HABIT termediate 3 = Open				
V	R1 2	R2 R3	R4		
V 1	N1 0	K2 K3			


2. LIGH	T SPROUT CHARACTERISTICS: (continued)			alle a co	
	LIGHT SPROUT TIP: PUBESCENCE 1 = Absent 2 = Weak 3 = Medium	4 = Strong	5 = Very Strong	20000(268
	V R1	R2	R3	R4	
	LIGHT SPROUT TIP ANTHOCYANIN COLO 1 = Green 2 = Red-violet 3 = Blue-		ther(describe)		_
	v 3 R1	R2	R3	R4	
•	LIGHT SPROUT TIP: INTENSITY OF ANTHO 1 = Absent 2 = Weak 3 = Medium	OCANIN COLORA 4 = Strong	ATION (IF PRESENT) 5 = Very Strong		
	V 5 R1 1	R2	R3	R4	
	LIGHT SPROUT ROOT INITIALS: FREQUE! 1 = Short	NCY			
	V 2 R1 2/2.5	R2	R3	R4	
3. PLAN	NT CHARACTERISTICS:				
	GROWTH HABIT: (See Figure 2)				
		ct (30-45° with gro	ound) 7 = Spreading		
	V R1	R2	R3	R4	
	TYPE: 1 = Stem (foliage open, stems clearly visible)	2 = Intermedia	te 3 = Leaf (Foliage cl	osed, stems hardly visible)	
	V R1	R2	R3	R4	
	MATURITY: Days after planting (DAP) at vis	ne senescence			
	V R1	R2	R3	R4	
	PLANTING DATE:				
	V R1	F	R2	R3	R4
	*REGIONAL AREA: 1 = Pacific North West (WA, OR, ID, CO, CA) 4 = Mid-Atlantic Erect (VI, NC, SC, South NJ, F) 7 = Europe 8 = England		Central (ND, WI, MI, MN, C (LA, TX, AZ, NE) a 10 = Brazil	0H) 3 = North East (ME, N 6 = Canada 11 = Other	Y, PA, NJ, MD, MA, RI,)
	V R1		R2	R3	R4
	MATURITY CLASS: 1 = Very Early (<100 DAP) 2 = Early (100-110) DAP) 3 = Mid-s	eason (111-120 DAP) 4 =	Late (121-130 DAP) 5 = Ve	y Late (>130 DAP).


R1

R4

Figure 1: Growth Habit








5 = Semi-erect $30-45^{\circ}$ with ground

7 = Spreading <30° with ground


Figure 2: Leaf Silhouette







3 = Medium



21

Figure 3: Terminal Leaflet Shape / Primary Leaflet Shape



1=Narrowly
Ovate



2=Medium Ovate



3=Broadly Ovate



4=Lanceolate



5=Elliptical



6=Obovate



7=Oblong

Figure 4: Terminal Leaflet Shape of Tip / Primary Leaflet Shape of Tip




Figure 5: Terminal Leaflet Shape of Base / Primary Leaflet Shape of Base






Figure 6: Corolla Shape











rotate i << b



semi-stellate I = b



pentagonal l < b



very rotate

Figure 7: Anther Shape




Figure 8: Stigma Shape




Figure 9: Distribution of Secondary Tuber Color




Figure 10: Tuber Shape




Figure 11: Leaf Dissection




Figure: 12 Stem Wings




Figure 13: Leaf Stipules:




EXHIBIT D. Additional Description of the Variety


- 1) Isozyme fingerprint of FL 1900, with reference to methodology. Comparison of fingerprint of FL 1900 with that of Atlantic, showing distinct patterns for each variety (Exhibit D-1).
- 2) Summary of 100-tuber sample of tuber dimensions of FL 1900 compared to 100 tubers of Atlantic


EXHIBIT 5.1

Isozy	me electro	phoresis	Isozyme electrophoresis fingerprints of FL 1900	of FL 19	00 com	pared to	compared to Atlantic							
Variety		Mdh-2	Mdh-1 Mdh-2 6-Pgdh-3 Idh-1 Pgi-1 Aps-1 Got-1 Got-2 Pgm-1 Pgm-2 Dia-1 Prx-1 Prx-3 Adh-1	ldh-1	Pgi-1	Aps-1	Got-1	Got-2	Pgm-1	Pgm-2	Dia-1	Prx-1	Prx-3	Adh-1
								:						
Atlantic	2223	2223	1122	1112	2222	1112 2222 1111 4444	4444	3555	3555 1112	2223	2223 1112 1144	1144	i	2222
FL1900	2244	2222	1122	1	2222	f	3344	3335	1223	2233	1	ı	1112	;

Source of Data: Dr. David Douches, Michigan State University, 1995

Procedures and allelic designations used are according to Douches, D.S. and K. Ludlam. 1991. Electrophoretic Characterization of North American Potato Cultivars. Am. Potato J. 68:767-780.





REPRODUCE LOCALLY. Include form num		reproductions.	FORM APPROVED - OMB No. 0581-00
U.S. DEPARTMENT OF AGE AGRICULTURAL MARKETIN EXHIBIT E		Application is required in order to certificate is to be issued (7 U.S. confidential until the certificate is	
STATEMENT OF THE BASIS	OF OWNERSHIP		
1. NAME OF APPLICANT(S)		2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBE	
Frito-Lay North America, Inc.	BPA SIZION	RD 78-91-5	FL 1900
4. ADDRESS (Street and No., or R.F.D. No., City, S	tate, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)
		972-334-3822	972-334-5965
7701 Legacy Drive	•	7. PVPO NUMBER	
Plano, Texas 75024	BOA S/z/as	200000268, 'FL 1900) *
8. Does the applicant own all rights to the v 9. Is the applicant (individual or company)			
			- [A] [
10. Is the applicant the original owner?	YES	X NO If no, please answer	one of the following:
a. If the original rights to variety were or b. If the original rights to variety were	X YES	NO If no, give name of co	puntry
	YES	NO If no, give name of co	
11. Additional explanation on ownership (If	needed, use the reverse fo	or extra space);	
The original owner (the obligations under his e pursuant to contractual	employment contr	act to Frito-Lay Inc	which in turn
PLEASE NOTE:			
Plant variety protection can only be afforded	d to the owners (not license	ees) who meet the following criteria	a:
If the rights to the variety are owned by the national of a country which affords similar	ne original breeder, that per protection to nationals of	rson must be a U.S. national, natio the U.S. for the same genus and s	onal of a UPOV member country, or species.
If the rights to the variety are owned by tr nationals of a UPOV member country, or genus and species.	ne company which employs owned by nationals of a co	ed the original breeder(s), the com buntry which affords similar protect	pany must be U.S. based, owned by tion to nationals of the U.S. for the same
3. If the applicant is an owner who is not the	original owner, both the o	riginal owner and the applicant mu	ust meet one of the above criteria.
The original breeder/owner may be the indiv Act for definitions.	ridual or company who dire	ected the final breeding. See Sect	ion 41(a)(2) ot the Plant Variety Protection
According to the Paperwork Reduction Act of 1995, an age control number. The valid OMB control number for this infresponse, including the time for reviewing the instructions,	omiation collection is USX1-11055	the time required to complete this information	collection is actimated to average 6 minutes nor

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

EXHIBIT E. Statement of the Applicant's Ownership

The variety FL 1900 for which Plant Variety Protection is hereby sought was developed by a breeder employed by Recot, Inc., who has assigned all rights to inventions and discoveries made by him to Recot, Inc., with no ownership rights of any kind retained by the breeder.

EXHIBIT F. Deposit Statement

Upon issuance of the Plant Variety Protection Certificate for FL 1900, applicant will deposit tissue culture plantlets of this variety in a public repository.

Form Approved OMB NO 0581-0055

REPRODUCE LOCALLY, Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT F DECLARATION REGARDING DEPOSIT

NAME OF OWNER (S) Frito-Lay North America, Inc.	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country) 7701 LEGACY DRIVE PLANO, TX 15024	TEMPORARY OR EXPERIMENTAL DESIGNATION 1991 78.05 VARIETY NAME FL 1900
NAME OF OWNER REPRESENTATIVE (S) Jondle & Associates, PC	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country) 858 HAPPY CANYON RD CASTLE ROCK, CO 80108	Pronumer 00268

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.